

Package: beers (via r-universe)

August 26, 2024

Title Calculate Beers Ordinary Interpolation

Version 0.1.0

Description Calculate Beers Interpolation (both Ordinary and Modified method); a six-term formula which minimizes the fifth differences of the interpolated results. See p877 of The Methods and Materials of Demography, Volume 2, referring to Beers, The Record of the American Institute of Actuaries, 34, Part I (69):59-60, June 1945.

License MIT + file LICENSE

Encoding UTF-8

LazyData true

RoxygenNote 6.0.1

Suggests knitr, rmarkdown, testthat

VignetteBuilder knitr

Repository <https://mrc-ide.r-universe.dev>

RemoteUrl <https://github.com/mrc-ide/beers>

RemoteRef master

RemoteSha 051aa0c40bddbf7d51210679708946db5a5f6d70

Contents

beers	2
beers_int_ordinary	2
beers_sub_ordinary	3

Index	4
--------------	----------

beers	<i>beers: A package for computing Beers ordinary and modified interpolation</i>
-------	---

Description

beers: A package for computing Beers ordinary and modified interpolation

beers_int_ordinary	<i>Beers interpolation, using ordinary or modified method.</i>
--------------------	--

Description

Create 4 interpolated points between each pair of given values. With the ordinary method, the given data points given will be included unchanged in the interpolated list. With the modified method, some smoothing occurs, and only the first and last points given are guaranteed to appear unchanged in the interpolated list.

Usage

```
beers_int_ordinary(points)
```

```
beers_int_modified(points)
```

Arguments

points A list of at least 6 numbers, eg, populations over time.

Value

A list with four extra interpolated points between each pair of given points.

Examples

```
# Interpolate population of UK (1950, 1955, 1960, 1965, 1970, 1975) to yearly points  
beers_int_ordinary(c(50616014, 51123707, 52433157, 54303107, 55634935, 56211947))  
beers_int_modified(c(50616014, 51123707, 52433157, 54303107, 55634935, 56211947))
```

beers_sub_ordinary *Beers subdivision, using ordinary or modified method.*

Description

With the ordinary method, each set of 5 subdivided values will always sum to the original data point given. With the modified method, some smoothing occurs, and this property is only true for the first and last given data point.

Usage

```
beers_sub_ordinary(points)
```

```
beers_sub_modified(points)
```

Arguments

points A list of at least 5 numbers, eg, populations by 5-year age-band.

Value

For subdivision: a list 5 times as long as the original, with each point subdivided into 5.

Examples

```
# Subdivide population of UK (2015), ages 0-4, 5-9, 10-14, 15-19, 20-24 into single years.
```

```
beers_sub_ordinary(c(4042918, 3927745, 3529200, 3779712, 4174572))
```

```
beers_sub_modified(c(4042918, 3927745, 3529200, 3779712, 4174572))
```

Index

beers, [2](#)
beers-package (beers), [2](#)
beers_int_modified
 (beers_int_ordinary), [2](#)
beers_int_ordinary, [2](#)
beers_sub_modified
 (beers_sub_ordinary), [3](#)
beers_sub_ordinary, [3](#)